

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

(10) International Publication Number
WO 02/45727 A1

WO 02/45727 A1

2001/76749	5 December 2001 (05.12.2001)	KR
2001/76750	5 December 2001 (05.12.2001)	KR
2001/76751	5 December 2001 (05.12.2001)	KR
2001/76752	5 December 2001 (05.12.2001)	KR

2001/76751	5 December 2001 (05.12.2001)	KR
2001/76752	5 December 2001 (05.12.2001)	KR

(71) Applicant (for all designated States except US): PLBIO CO., LTD. [KR/KR]; 570-1, Sillimdong, Kwanak-gu, Seoul 151-877 (KR).

English

English

(75) **Inventors/Applicants (for US only):** LEE, Yeonhee [KR/KR]; Pacific House B-203, 129-2, Anam-5-ga, Sungbuk-gu, Seoul 136-075 (KR). PAEK, Kyungsoo [KR/KR]; Pacific House B-203, 129-2, Anam-5-ga, Sungbuk-gu, Seoul 136-075 (KR).









2000/74602	8 December 2000 (08.12.2000)	KR
2001/2577	17 January 2001 (17.01.2001)	KR
2001/2578	17 January 2001 (17.01.2001)	KR
2001/2579	17 January 2001 (17.01.2001)	KR
2001/8373	20 February 2001 (20.02.2001)	KR
2001/40135	5 July 2001 (05.07.2001)	KR
2001/40136	5 July 2001 (05.07.2001)	KR
2001/40137	5 July 2001 (05.07.2001)	KR
2001/40138	5 July 2001 (05.07.2001)	KR
2001/40139	5 July 2001 (05.07.2001)	KR
2001/40134	5 July 2001 (05.07.2001)	KR
PCT/KR01/01286	27 July 2001 (27.07.2001)	KR
2001/76748	5 December 2001 (05.12.2001)	KR

(74) Agent: **YOUME PATENT AND LAW FIRM**; Teheran Bldg., 825-33, Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,

[Continued on next page]

(54) Title: LACTIC ACID BACTERIA INHIBITING ADHESION OF *HELICOBACTER PYLORI* TO GASTRIC MUCOSA

Control	Competitive Lipid Binding Assay		
Lipid	PL9001	PL9002	PL9003
			
<i>Helicobacter pylori</i>	PL9004	PL9005	PL9006
			

(57) Abstract: The present invention relates to lactic acid bacteria capable of inhibiting activities of *Helicobacter pylori*, and more particularly, to lactic acid bacteria having an inhibitory activity on growth of *Helicobacter pylori* causing stomach ulcer and adhesion to the gastric mucosa, wherein the lactic acid bacteria are selected from the group consisting of coprophilus PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus* PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251): The lactic acid bacteria of the invention can be used as antiulcer drug, food additives, drugs for the prevention or treatment of *Helicobacter pylori*, drugs against bacteria that cause food poisoning, or drugs for the prevention or treatment of infectious bacteria.

WO 02/45727 A1



MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**LACTIC ACID BACTERIA INHIBITING ADHESION OF *HELICOBACTER*
PYLORI TO GASTRIC MUCOSA**

BACKGROUND OF THE INVENTION

(a) Field of the Invention

5 The present invention relates to lactic acid bacteria capable of inhibiting activities of *Helicobacter pylori*, and more particularly, to lactic acid bacteria having an inhibitory activity on growth of *Helicobacter pylori* causing stomach ulcer and adhesion to the gastric mucosa.

(b) Description of the Related Art

10 *Helicobacter pylori* is a gram-negative bacterium. In 1983, Drs. Barry Marshall and Robin Warren of Australia discovered the bacteria in the stomach lining of patients with chronic gastritis and peptic ulcers. *Helicobacter* was initially called *Campylobacter pyloridis*, and was later named as *Helicobacter pylori* from its morphology in the body and the name
15 of its habitat, the *pylorus*.

Helicobacter pylori infection is an important environmental factor in the pathogenesis of peptic ulcer disease, gastric carcinoma, and gastric lymphoma (Dubois, A. 1995. Spiral bacteria in the human stomach: the gastric *Helicobacters*. *Emerg. Infect. Dis.* 1: 79-85; Slomiany, B. L. and A.
20 Slomiany. 1992. Mechanism of *Helicobacter pylori* pathogenesis: focus on mucus. *J. Clin. Gastroenterol.* 14 Suppl 1 S114-21). Once *Helicobacter*

pylori infects, it remains for several decades and is not eliminated naturally, and thus *Helicobacter pylori* is a major cause of chronic gastritis.

Helicobacter pylori infects through the intake of food, and it attaches to the gastric mucosa and the duodenal mucosa. A pathogenic factor of *Helicobacter pylori* is a urease that makes it for surviving in the highly acidic condition of a stomach, a flagellum for maintaining mobility, and the outer membrane protein having adherence to the gastric mucosa.

In its attachment to the human gastric epithelium, *Helicobacter pylori* binds with the same antigens identified as that of red blood cells (Alkout, A. M., C. C. Blackwell, D. M. Weir, I. R. Poxton, R. A. Elton, W. Luman, and K. Palmer. 1997. *Gastroenterol.* 112: 1179-1187; Boren, T., P. Flak, K. A. Roth, G. Larson, and S. Normark. 1993. *Science* 262: 1892-1895; Clyne, M. and B. Drumm. 1997. *Gastroenterol.* 113: 72-80). Antigens, like the Lewis antigen isolated in human blood type O, are expressed in the gastric mucosa. Therefore, gastritis breaks out mostly in blood-type O patients. (Heneghan, M. A., A. P. Moran, K. M. Feeley, E. L. Egan, J. Goulding J, C. E. Connolly, and C. F. McCarthy. 1998. *FEMS Immunol. Med. Microbiol.* 20: 257-266; Kobayashi, K., J. Sakamoto, Y. Yamamura, T. Kito, H. Inagaki, T. Watanabe, and H. Nakazato. 1991. *Nippon Geka Gakkai Zasshi* 92: 813-819; Kobayashi, K., J. Sakamoto, Y. Kito, Y. Yamamura, T. Koshikawa, M. Fugita, T. Watanabe, and H. Nakazato. 1993. *Am. J. Gastroenterol.* 88: 919-924).

For the treatment of *Helicobacter pylori* infection, antibiotics, inhibitor

of proton pumping, and antacid have been used. Because *Helicobacter* is difficult to culture with a large scale, a vaccine using the whole bacterium has been developed unsuccessfully. The method using antibiotics has disadvantage in that the *Helicobacter pylori* become resistant to the antibiotics and can be infected again. Method using antacid that suppresses secretion of gastric acid is not a basic solution. In addition, although a vaccine using urease has been developed, it is not effective. In the future, the development of a vaccine against *Helicobacter pylori* will be difficult, due to its complex culture conditions, which make it difficult to determine an active area for a vaccine.

SUMMARY OF THE INVENTION

The present invention is designed by the necessities of the prior art, and it is an object of the present invention to provide bacteria that are capable of suppressing stomach ulcers.

It is another object to provide bacteria that are capable of inhibiting the adhesion of *Helicobacter pylori* to the gastric mucosa.

It is still another object to provide bacteria that inhibit the activity of *Helicobacter pylori*.

It is still another object to provide bacteria that inhibit the growth of *Helicobacter pylori*.

It is still another object to provide bacteria that inhibit the growth of bacteria that cause food poisoning.

It is still another object to provide bacteria that inhibit the growth of anaerobic bacteria or bacteria that cause acne.

It is still another object to provide bacteria that improve immunity.

In order to achieve these objects, the present invention provides a
5 bacterium for inhibiting the adherence of *Helicobacter pylori* to the gastric mucosa, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.

The present invention provides a bacterium for inhibiting the growth
10 of *Helicobacter pylori*, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.

The present invention provides a composition for inhibiting the growth of bacteria, wherein the composition comprises at least one bacteria
15 selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.

The present invention provides a cosmetic composition comprising at least one bacteria selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus*
20 *fermentum*, and a culture filtrate of bacteria thereof.

The present invention provides a food additive comprising at least one bacteria selected from the group consisting of *Lactobacillus coprophilus*,

Enterococcus durans, *Streptococcus faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

The present invention provides a food prepared by fermenting at least one bacteria selected from the group consisting of *Lactobacillus*
5 *coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.

The present invention provides a immunological enhancement composition comprising at least one bacteria selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus*
10 *faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

The present invention provides a composition for curing intestinal disorders comprising at least one bacteria selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

15

BRIEF DESCRIPTION OF THE DLIVEINGS

FIG. 1 is a slide showing *Helicobacter pylori* attachment to glycolipids isolated from red blood cells, wherein the amount of adhered *Helicobacter pylori* is proportional to the amount of glycolipids (0 ug, 0.14 ug, 1.4 ug, 14 ug, 70 ug, 140 ug, 280 ug);

20

FIG. 2 is TLC plate that shows inhibition of *Helicobacter pylori* adhesion to glycolipids by PL bacteria;

FIG. 3 is a photograph showing inhibition of *Helicobacter pylori*

adhesion to the gastric mucosa by PL bacteria;

FIG. 4 is a photograph showing PL bacteria adhered to an MKN-45 cell;

FIG. 5 shows that materials produced from PL bacteria suppress the
5 growth of *Helicobacter pylori*;

FIG. 6 is a graph showing the diameter of the growth-inhibited-zone for *Helicobacter pylori*;

FIG. 7 is a graph showing the growth-inhibited degree of *Helicobacter pylori* by a culture filtrate of PL bacteria;

10 FIG. 8 is a graph showing growth inhibition of bacteria that cause food poisoning by *Lactobacillus coprophilus* PL 9001;

FIG. 9 is a graph showing growth inhibition of bacteria that cause food poisoning by *Enterococcus durans* PL 9002;

FIG. 10 is a graph showing growth inhibition of bacteria that cause
15 food poisoning by *Streptococcus faecalis* PL 9003;

FIG. 11 is a graph showing growth inhibition of bacteria that cause food poisoning by *Lactobacillus coprophilus* PL 9004;

FIG. 12 is a graph showing growth inhibition of bacteria that cause food poisoning by *Lactobacillus fermentum* PL 9005;

20 FIG. 13 is a graph showing growth inhibition of bacteria that cause food poisoning by *Lactobacillus fermentum* PL 9006;

FIG. 14 is a graph showing growth inhibition of bacteria that cause acne by PL bacteria;

FIG. 15 is a graph showing an immunological enhancement effect of PL bacteria; and

5 FIG. 16 is photograph showing the adhered PL bacteria to CaCo-2 cell lines.

DETAILED DESCRIPTION OF THE PRESENT INVENTION

The present invention provides bacteria that inhibit the adhesion of *Helicobacter pylori* to the gastric mucosa. The bacteria are lactic acid
10 bacteria and preferably *Lactobacillus sp.* or *Enterococcus sp.* More preferably, the bacteria are *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*, and most preferably, *Lactobacillus coprophilus* PL 9001, *Enterococcus durans* PL 9002, *Streptococcus faecalis* PL 9003, *Lactobacillus coprophilus* PL 9004,
15 *Lactobacillus fermentum* PL 9005, and *Lactobacillus fermentum* PL 9006. The *Streptococcus faecalis* is identical to *Enterococcus faecalis*: its name has been changed to *Streptococcus faecalis* PL 9003.

The PL bacteria (*Lactobacillus coprophilus* PL 9001, *Enterococcus durans* PL 9002, *Streptococcus faecalis* PL 9003, *Lactobacillus coprophilus*
20 PL 9004, *Lactobacillus fermentum* PL 9005, and *Lactobacillus fermentum* PL 9006) have been internationally deposited in the Korean Culture Center of Microorganisms and given accession numbers. The accession numbers

given by the international depositary authority are KCCM-10245 for *Lactobacillus coprophilus* PL 9001, KCCM-10246 for *Enterococcus durans* PL 9002, KCCM-10247 for *Streptococcus faecalis* PL 9003, KCCM-10248 for *Lactobacillus coprophilus* PL 9004, KCCM-10250 for *Lactobacillus fermentum* PL 9005, and KCCM-10251 for *Lactobacillus fermentum* PL 6.

The present invention provides an antiulcer drug. The antiulcer drug contains *Lactobacillus sp.* or *Enterococcus sp.*, and more preferably it contains the PL bacteria. The *Lactobacillus sp.* or *Enterococcus sp.* have inhibitory activities for *Helicobacter pylori* adhesion to the gastric mucosa as well as for *Helicobacter pylori* growth. These activities are confirmed with *Lactobacillus coprophilus* PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus* PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251).

Also, the PL bacteria produce materials that inhibit the growth of *Helicobacter pylori* and adhesion thereof to the gastric mucosa, and thus, a culture filtrate of the PL bacteria can be used for the anti-*Helicobacter* materials. The PL bacteria have antibiotics-resistant activity, acid-resistant activity, and bile-resistant activity, and are stable *in vivo*, and the PL bacteria can be provided as live bacteria, dehydrated bacteria, and non-viable bacteria.

The antiulcer drug of the present invention can be administered via oral or parenteral routes. Preferably, the drug would be administered orally or by injection, but most preferably, the drug would be administered orally. The antiulcer can be prepared as a single compound drug or a complex drug, and the complex drug can contain more pharmaceutically acceptable material. It is reasonable that the blending ratio of the PL bacteria is controlled according to the type of drug.

The culture filtrate of PL bacteria is a solution without bacteria that is prepared by filtering or centrifuging culture fluid, and more preferably, the culture filtrate is a supernatant phase prepared by inoculating the PL bacteria in a MRS liquid broth, culturing at 37°C for 16-48 hr, and centrifuging.

In addition, *Lactobacillus sp.* and *Enterococcus sp.* can be applied to inhibit the activity of other bacteria. More preferably, the PL bacteria (PL9001, PL9002, PL9003, PL9004, PL9005, and PL9006), fragmented cell walls of the PL bacteria, or the culture filtrate of the PL bacteria can be used to inhibit the growth of bacteria that cause food poisoning, anaerobic bacteria, bacteria that cause acne, and so on.

The anaerobic bacteria are general disease-causing germs. The anaerobic bacteria are preferably tetanus bacteria, gas gangrene bacteria, and *Clostridium sp.*, and more preferably, *Clostridium perfringens*.

The bacteria that cause food poisoning are bacteria such as a listeriosis, dysentery, diarrhea, O157, and a bacteria that mediates food

poisoning.

PL bacteria of the present invention have activity for immunological enhancement, they have adhesion capability to gastric and intestinal cells, and they can be used to improve health. The PL bacteria can be administered to human beings or to animals for therapeutic or prophylactic treatment of the stomach and particularly the pylorus, for inhibiting growth of harmful bacteria, and for curing intestinal disorders. The PL bacteria and their culture filtrate are capable of displacing pathogenic bacteria from intestinal or gastric cells to which they adhere.

10 The present invention provides a composition including the PL bacteria. The composition can be used for inhibiting biological activity of *Helicobacter pylori* and bacteria, for immunity improving, or as a medicine for intestinal disorders, and it can be applied to feed, feed additives, food, food additives, medicine, or cosmetic compositions.

15 The composition contains the cell walls prepared by grinding the live, non-viable, and dehydrated PL bacteria or a culture filtrate thereof as an effective agent, and it further contains excipients or a carrier. The composition containing the excipients or carrier includes 0.001 to 90 wt% of PL bacteria, but it is not limited to that. The PL bacteria of the present invention have undergone a toxicity test, which has confirmed that the PL bacteria are safe *in vivo*.

The feed or feed additives include 0.001 to 90 wt% of PL bacteria or

a culture filtrate thereof, but its amount is not limited to that. The feed or feed additives are prepared as a drink or a solid product.

The food or food additives are preferably yogurt, weaning diet, dairy goods, cheese, Kimchi, drinks, or other food, and they are prepared by
5 fermenting with the PL bacteria of the present invention as a seed.

The medicine may be used by itself or it may be included with more than one pharmaceutical composition. The medicine can include more than one kind of pharmaceutical diluent selected from the group consisting of saline, buffered saline, dextrose, water, glycerol, and ethanol, but the diluent
10 is not limited. Appropriate diluents are listed in the written text of Remington's Pharmaceutical Science (Mack Publishing Co, Easton PA).

The medicine may be applied differently according to the purpose of dosing and diseases. It should be understood that the amount of the active ingredient actually administered ought to be determined in light of various
15 relevant factors including the condition to be treated, the severity of the patient's symptoms, co-administration with other drugs (e.g., chemotherapeutic agents), the age, sex, and body weight of the individual patient, food consumed, dosing time, the chosen route of administration, and the ratio of the composition.

20 A daily dose of the medicine is preferable from about 1 to 800 mg, and most preferably 40 to 400 mg. In general, 0.1 to 20 mg/kg, and preferably 0.2 to 10 mg/kg of the medicine can be administrated in a single

or in 1-3 divided doses per day, even though the exact dose and route of administration are adjusted to the type and severity of disease.

The medicine containing PL bacteria or a culture filtrate thereof can be administered via oral or parenteral routes. Parenteral dosing means the
5 administration of a drug through a route other than oral, which includes rectal, intravenous, intraperitoneal and intramuscular, intra-arterial, transdermal, nasal, inhalation, ocular, and subcutaneous introduction.

Pharmaceutical formulations may be prepared in any form, such as an oral dosage form, an injectable solution, suppository or a topical
10 preparation. The formulation is preferably prepared for oral and injectable administration (true solution, suspension, or emulsion), and most preferably in oral form such as tablet, capsule, soft capsule, aqueous medicine, pill, granules, and the like.

In preparing the formulation, the PL bacteria or a culture filtrate are
15 filled in the soft capsule without any excipients, or formed as an appropriate formulation after mixing or diluting with a carrier. Examples of suitable carriers are starches, water, saline, Ringer's solution, dextrose, and any ingredients described in previous reports (e.g. Remington's Pharmaceutical Science, Mack Publishing Co., Easton PA).

20 The cosmetic composition contains 0.001 to 20 wt% of PL bacteria or a culture filtrate thereof, but the ratio is not limited to that. The cosmetic composition further contains a general material added to cosmetics such as

oil, water, detergent, moisturizer, alcohol, thickener, chelate compound, pigment, preservative, and perfume, in appropriate amounts. The cosmetic compound can be formulated as a toner, lotion, cream, essence, pack, powder, ointment, suspension, emulsion, or spray. The cosmetic
5 composition may be used for prevention and treatment of bacteria, such as those causing acne, and it is preferably for external application.

The present invention provides a material for inhibiting the growth of harmful bacteria. The material for inhibiting the growth of harmful bacteria is able to suppress the growth of *Helicobacter pylori*, anaerobic bacteria,
10 bacteria that cause acne, or bacteria that cause food poisoning, and it is prepared as a culture filtrate in a bacteria culture. Thus, the material for inhibiting the growth of harmful bacteria of *Lactobacillus* sp. or *Enterococcus* sp., can also be used for the treatment and prevention of bacteria.

As mentioned above, *Lactobacillus coprophilus* PL 9001,
15 *Enterococcus durans* PL 9002, *Streptococcus faecalis* PL 9003, *Lactobacillus coprophilus* PL 9004, *Lactobacillus fermentum* PL 9005, and *Lactobacillus fermentum* PL 9006 of the present invention suppress the growth and adhesion of *Helicobacter pylori*. Also, PL bacteria, fragments of PL bacteria, or a culture filtrate of PL bacteria can be used for the treatment
20 and prevention of bacterial infection. PL bacteria and a culture filtrate of PL bacteria inhibit the growth of bacteria that cause food poisoning, bacteria that cause acne, and anaerobic bacteria, as well as enhance immunity effects,

and thereby PL bacteria and a culture filtrate of PL bacteria can be used for treatment and prevention of bacterial manifestations.

The present invention will be explained in more detail with reference to the following Examples. However, the following Examples are to
5 illustrate the present invention and the present invention is not limited to them.

EXAMPLE 1: Isolation of PL bacteria

MRS broth (Difco, bacto proteose peptone No.3 10 g, bacto beef extract 10 g, bacto yeast extract 5 g, bacto dextrose 20 g, polysorbate 80 g,
10 ammonium citrate 2 g, sodium acetate 5 g, magnesium sulfate 0.1 g, manganese sulfate 0.05 g, and dipotassium phosphate 2 g /L) was mixed with 0.002% of bromophenol blue so that a MRS+BPB medium was prepared.

Kimchi samples were diluted with a 10 X pepton solution. 0.1 mL of
15 the diluted solution was inoculated into the MRS+BPB medium and spread. The feces from an infant were picked by a cotton swab and inoculated into the MRS+BPB medium. After incubation for 3-4 days in an incubator (25°C), lactic acid bacteria were observed and isolated according to the colony form.

In order to distinguish a group of lactic acid bacteria, a MRS solid
20 medium was added to BPB which displays a yellow color at pH 3.0 and a purple color at pH 4.6, and the lactic acid bacteria were classified according to coloration by the BPB. The coloration was determined by producing-

degrees of lactic acid, pH-resistance, and the length of life. *P. acidolactic* and *S. faecalis* are normal lactic acid bacterium, *L. mesenteroides* and *L. brevis* are abnormal lactic acid bacterium, and *L. plantarum* is a random lactic acid bacterium. *S. faecalis* forms a white colony where its medium
5 changes to light yellow. *L. mesenteroides* ferments abnormal lactic acid and produces a low level of lactic acid, and the colony is deep blue as a whole and has a small size without a ring. *Lactobacillus sp.* has a light blue ring, being deep blue in the center, or a generally white color, and the size of the colony is large. *P. acidolactic* and *L. mesenteroides* express a deep-
10 blue because of a short lifetime and low pH. For example, *L. mesenteroides* cannot grow below pH 4.8.

All bacteria of the present invention formed a white colony of over 0.3 mm (diameter) and were classified as *Lactobacillus sp.*

Bacteria were isolated from a single colony, according to Bergy's
15 manual of systematic bacteriology, by exhibition of morphological and biochemical properties. After performed a gram stain and a catalase-reaction, the bacteria were analyzed in an API system (La Balme-les-Grottes, France). After the colony was picked with a cotton swab, it was floated with 2 ml of distilled water. The floating solution was added to an API 50 CHL
20 medium and mixed. 200 ul of the mixture were inoculated into each of 50 tubes, and each mixture was covered with mineral oil and incubated at 37°C for 48 hr.

The fermentation pattern of 49 kinds of saccharides were analyzed with API 50 CH and API 50 CHL, the resulting data were fed into the ATB identification computer system (bio Merieux France), and then the bacterium was identified. The results of the first bacterium among the isolated 5 microorganisms are presented in Table 1.

(Table 1)

Strip No.1 tube/substrate	Strip No.2 tube/substrate	Strip No.3 tube/substrate	Strip No.4 tube/substrate	Strip No.5 tube/substrate
- Control	- Galactose	- D-Mannoside	- Melibiose	- D-Turanose
- Glycerol	+ D-Glucose	- D-Glucoside	+ Saccharose	- D-Lyxose
- Erythritol	+ D-Fructose	+ Glucosamine	- Trehalose	- D-Tagatose.
- D-Arabinose	+ D-Mannose	+ Amygdaline	- Inuline	- D-Fucose
- L-Arabinose	- L-Sorbose	+ Arbutine	- Melezitose	- L-Fucose
- Ribose	- Rhamnose	+ Esculine	- D-Raffinose	- D-Arabitol
+ D-Xylose	- Dulcitol	+ Salicine	- Amidon	- L-Arabitol
- L-Xylose	- Inositol	+ Cellobiose	- Glycogene	+ Gluconate
- Adonitol	- Mannitol	+ Maltose	- Xylitol	- 2-Gluconate
- Xyloside	- Sorbitol	- Lactose	+ Gentiobiose	- 5-Gluconate

The first bacterium has a 99.9% similarity to *Lactobacillus coprophilus* and a 0.1% similarity to *Lacto. brevis*. 16S rRNA of the first bacterium was sequenced with a 16S rRNA gene kit (Perkin Elmer Applied Biosystem) and the sequence is shown as Sequence No. 1. The base sequence of Sequence No. 1 is identical to the 16S rRNA base sequence of *Lactobacillus coprophilus* as found from a BLAST search (<http://www.ncbi.nlm.nih.gov/blast>). Other names of *Lactobacillus coprophilus* are *Weissella confusa* and *Lactobacillus confusus*. The 15 bacterium was deposited under *Lactobacillus* 9001 (KFCC-11240) in the Korean Federation of Culture Collections, and internationally deposited

under *Lactobacillus coprophilus* PL 9001(KCCM-10245) in the Korean Culture Center of Microorganisms.

The second bacterium was characterized with an API STREP Kit and the results are presented in Table 2.

5 (Table 2)

VP	+	ARA	+
HIP	-	MAN	+
ESC	+	SOR	-
PYRA	+	LAC	-
AGAL	-	TRE	-
β GUR	-	INU	-
β GAL	-	RAF	-
PAL	-	AMD	-
LAP	-	GLYG	-
ADH	+	β - HEM	-
RIB	+		

For the non-classified second bacterium, the 16S rRNA base sequence was analyzed. The 16S rRNA base sequence of the second bacterium is Sequence No. 2, and it is identical to that of *Enterococcus durans*. According to the sequencing result, the second bacterium was
 10 called *Enterococcus durans* PL 9002, and it was deposited under *Enterococcus durans* PL9002 (KFCC-11241) in the Korean Federation of Culture Collections, and internationally deposited under *Enterococcus durans* PL 9002 (KCCM-10246) in the Korean Culture Center of Microorganisms.

15 The results of the third bacterium characterization are presented in Table 3.

(Table 3)

VP	+	ARA	-
----	---	-----	---

HIP	-	MAN	+
ESC	+	SOR	+
PYRA	+	LAC	+
AGAL	-	TRE	+
β GUR	-	INU	-
β GAL	-	RAF	-
PAL	-	AMD	+
LAP	+	GLYG	-
ADH	+	β - HEM	-
RIB	+		

The third bacterium was identical to *Streptococcus faecalis* (*Enterococcus faecalis*) by 99.1%. The bacterium was deposited under *Streptococcus faecalis* 9003 (KFCC-11242) in the Korea Federation of Culture Collections, and internationally deposited under *Streptococcus faecalis* PL 9003(KCCM-10247) in the Korea Culture Center of Microorganisms.

The 16S rRNA base sequence of the third bacterium was analyzed as Sequence No. 3, and it was found to be identical to that of *Streptococcus faecalis* (*Enterococcus faecalis*).

10 The results of the fourth bacterium characterization are presented in Table 4.

(Table 4)

Strip No.1 tube/substrate	Strip No.2 tube/substrate	Strip No.3 tube/substrate	Strip No.4 tube/substrate	Strip No.5 tube/substrate
- Control	- Galactose	- D-Mannoside	- Melibiose	- D-Turanose
- Glycerol	+ D-Glucose	- D-Glucoside	+ Saccharose	- D-Lyxose
- Erythritol	+ D-Fructose	+ Glucosamine	- Trehalose	- D-Tagatose
- D-Arabinose	+ D-Mannose	+ Amygdaline	- Inuline	- D-Fucose
+ L-Arabinose	- L-Sorbose	- Arbutine	- Melezitose	- L-Fucose
- Ribose	- Rhamnose	+ Esculine	- D-Raffinose	- D-Arabitol
+ D-Xylose	- Dulcitol	+ Salicine	- Amidon	- L-Arabitol
- L-Xylose	- Inositol	+ Cellobiose	- Glycogene	+ Gluconate
- Adonitol	- Mannitol	+ Maltose	- Xylitol	- 2-Gluconate
- Xyloside	- Sorbitol	- Lactose	+ Gentiobiose	- 5-Gluconate

The fourth bacterium was identical to *Lactobacillus coprophilus* by 98.2% and to *Lacto. brevis* by 1.3%. Also, the 16S rRNA base sequence of the bacterium was analyzed as Sequence No. 4, and it was found to be identical to that of *Lactobacillus coprophilus* as a result of a BLAST search (http://www.ncbi.nlm.nih.gov/blast). The bacterium was deposited under *Lactobacillus coprophilus* PL 9004 (KFCC-11243) in the Korea Federation of Culture Collections, and internationally deposited under *Lactobacillus coprophilus* PL 9-4 (KCCM-10248) in the Korean Culture Center of Microorganisms

The results of the fifth bacterium characterization are presented in Table 5.

(Table 5)

Strip No.1 tube/substrate	Strip No.2 tube/substrate	Strip No.3 tube/substrate	Strip No.4 tube/substrate	Strip No.5 tube/substrate
- Control	+ Galactose	- D-Mannoside	+ Melibiose	- D-Turanose
- Glycerol	+ D-Glucose	- D-Glucoside	+ Saccharose	- D-Lyxose
- Erythritol	+ D-Fructose	+ Glucosamine	- Trehalose	- D-Tagatose
- D-Arabinose	+ D-Mannose	- Amygdaline	- Inuline	- D-Fucose
- L-Arabinose	- L-Sorbose	- Arbutine	- Melezitose	- L-Fucose
+ Ribose	- Rhamnose	- Esculine	+ D-Raffinose	- D-Arabitol
- D-Xylose	- Dulcitol	- Salicine	- Amidon	- L-Arabitol
- L-Xylose	- Inositol	- Cellobiose	- Glycogene	+ Gluconate
- Adonitol	- Mannitol	+ Maltose	- Xylitol	- 2-Gluconate
- Xyloside	- Sorbitol	- Lactose	- Gentiobiose	- 5-Gluconate

The fifth bacterium was identical to *Lactobacillus fermentum* by 93.2% and to *Leuconostoc lactis* by 6.7%. Also, the 16S rRNA base sequence of the bacterium was analyzed as Sequence No. 5, and it was found to be identical to that of *Lactobacillus fermentum* as a result of a

BLAST search (<http://www.ncbi.nlm.nih.gov/blast>). The bacterium was deposited under *Lactobacillus fermentum* PL 9005 (KCCM-10250) in the Korean Culture Center of Microorganisms.

The results of the sixth bacterium characterization are presented in

5 Table 6.

(Table 6)

Strip No.1 tube/substrate	Strip No.2 tube/substrate	Strip No.3 tube/substrate	Strip No.4 tube/substrate	Strip No.5 tube/substrate
- Control	+ Galactose	+ D-Mannoside	+ Melibiose	- D-Turanose
- Glycerol	+ D-Glucose	- D-Glucoside	+ Saccharose	- D-Lyxose
- Erythritol	+ D-Fructose	- Glucosamine	+ Trehalose	- D-Tagatose
- D-Arabinose	+ D-Mannose	- Amygdaline	- Inuline	- D-Fucose
- L-Arabinose	- L-Sorbose	- Arbutine	- Melezitose	- L-Fucose
+ Ribose	- Rhamnose	- Esculine	+ D-Raffinose	- D-Arabitol
- D-Xylose	- Dulcitol	- Salicine	- Amidon	- L-Arabitol
- L-Xylose	- Inositol	- Cellobiose	- Glycogene	- Gluconate
- Adonitol	- Mannitol	+ Maltose	- Xylitol	- 2-Gluconate
- Xyloside	- Sorbitol	+ Lactose	+ Gentiobiose	- 5-Gluconate

The sixth bacterium was identical to *Lactobacillus fermentum* by 94.4% and to *Lactobacillus lactis* by 5.4%. Also, the 16S rRNA base sequence of the bacterium was analyzed as Sequence No. 6, and it was
 10 found to be identical to that of *Lactobacillus fermentum* as a result of a BLAST search (<http://www.ncbi.nlm.nih.gov/blast>). The bacterium was deposited under *Lactobacillus fermentum* PL 9006 (KCCM-10251) in the Korean Culture Center of Microorganisms

EXAMPLE 2. Test for inhibiting adherence of *Helicobacter pylori* to

15

the gastric mucosa

(1) Preparation of bacteria

Helicobacter pylori (ATCC 43504) was cultured for 48 hr on a Brucella solid medium (Brucella broth, fungizone (2.5 g/ml amphotericin B) with Skirrow's supplement (0.16 mg/ml polymyxin B, 5 mg/ml vancomycin, 2.5 mg/ml trimethoprim)) and further supplemented with 10% horse serum, under a 5-10% CO₂ incubator. Cultured cells were collected by scraping, they were washed twice with a phosphate-buffered saline (PBS, pH 7.4), and then kept in a 10 mM Tris-Cl buffer at -20°C until used.

The bacteria prepared by EXAMPLE 1 is referred to as PL bacteria. The PL bacteria (*Lactobacillus coprophilus* PL 9001(KCCM-10245), *Enterococcus durans* PL 9002(KCCM-10246), *Streptococcus faecalis* PL 9003(KCCM-10247), *Lactobacillus coprophilus* PL 9004(KCCM-10248). *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251)) were grown for 24 hr at 37°C in an MRS broth, collected by centrifugation, pellet washed 2-3 times with PBS (pH 7.4), and then kept in 10 mM Tris-cl at -20°C until used.

(2) Extraction and purification of glycolipid

Glycolipid was isolated from human type O red blood cells (RBCs), as described in a report by Lee et. al. (Lee, Y., E. Shin, J. Lee, and J.H. Park. 1999. *J. Microb. Biotech.* 9: 794-797), with a slight modification. The human type O RBCs were dispersed in a minimum volume of water, repeated processed by freezing at -70°C and melting, until they were broken down. After the lysate was fixed, the supernatant was removed. The

remains were collected and extracted as mentioned below, and then mixed with a chloroform-methanol mixture (2:1, v/v), and a lower phase (containing lipid) was collected. The lipid layer material was dried in a rotary vacuum evaporator. The dried solid was dissolved in chloroform containing 2% methanol and loaded into a column (bed volume = 20 ml) that was equilibrated with silicic acid (H_2SiO_3). The fractions were collected by sequentially eluting with chloroform, acetone-methanol (3:1, v/v), and methanol. The methanol fraction was then dried in a rotary vacuum evaporator, dissolved in a minimum volume of methanol, and stored in microcentrifuge tube at -70°C .

(3) Competitive lipid binding assay

A competitive lipid binding assay (Lee, Y., E. Shin, J. Lee, and J.H. Park. 1999. *J. Microb. Biotech.* 9: 794-797) was performed with a slight modification. The extracted glycolipid (14 ug/5 ul) was spotted on a thin layer chromatography plate (TLC, Merck, Kieselgel60, EM Separations, Gibbstown, NJ, U.S.A). The plate was soaked in a 10 mM Tris buffer (pH 7.6) containing 3% gelatin for 2 hr at 37°C to prevent non-specific binding. The TLC plate was rinsed twice with the same buffer and a culture solution of PL bacteria ($\text{OD}_{600}=1$, CFU 2.4×10^8) was added. After 5 to 30 min, *H. pylori* (2.0×10^8 CFU) was added and the mixture was gently agitated for 2 hr at 37°C . In addition, except for the PL bacteria, the control plate was reacted with only *Helicobacter pylori* as described above, and agitated for 2

hr. The plate was rinsed 3 times with the same buffer for 10 min/time. The first antibody, rabbit antiserum raised against *H. pylori*, was added to the buffer (1: 600) and further incubated for 2 hr at room temperature with gentle shaking. After the plate was washed to remove the first antibody (1: 1000),
5 IgG, the second antibody conjugated with alkaline phosphatase, was added to the plate and the plate was incubated for 1 hr at room temperature. A chromogenic reaction was observed by adding BCIP/NBT (5-bromo-4-chloro-3-indolyl phosphate disodium salt/nitro blue tetrazolium chloride). Therefore, the chromogenic level showed the adherence degree of *Helicobacter pylori*
10 on glycolipid derived from red blood cells.

FIG. 1 is a TCL plate showing that with *Helicobacter pylori* bound to glycolipid extracted from red blood cells, the amounts of bound *H. pylori* increased linearly as the amount of spotted glycolipid (0 ug, 0.14 ug, 1.4 ug, 14 ug, 70 ug, 140 ug, 280 ug) increased.

15 FIG. 2 is TLC plate for observing whether or not *Lactobacillus* inhibits the adhesion of *Helicobacter pylori* on glycolipids. PL bacteria (*Lactobacillus coprophilus* PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002(KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus* PL 9004 (KCCM-10248), *Lactobacillus*
20 *fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251)) was reacted with *Helicobacter pylori*. As shown in FIG. 2, PL bacteria of the EXAMPLE inhibited the adherence of *Helicobacter*

pylori.

(4) Test for inhibiting adherence of *Helicobacter pylori* to the gastric mucosa

Bacteria having biological activity and non-viable bacteria prepared
5 by boiling at 75°C for 15 min were prepared. The adhesion assay was conducted with the surviving bacteria and the non-viable bacteria on MKN-45 (Human Gastric Adenocarcinoma).

MKN cells ($2.8-3.0 \times 10^5$) were added to each plate (diameter 30 mm) and cultured in a CO₂ incubator for 5-6 days until confluency. After cells
10 were confluent, cells were washed three times with PBS (phosphate-buffered saline), a new medium was added, and cultured further 30 min at 37°C. *Helicobacter pylori* (10^6 CFU) and PL bacteria (10^7 CFU) were added to these MKN cells and incubated for 1 hour. Unbound PL bacterial cells were removed by three times washing with PBS. Then cells were fixed in a fixing
15 solution (100 ml 3.5% formaldehyde, 16 g Na₂HPO₄, 4 g NaH₂PO₄·H₂O, D.W. to 1 liter) and kept for 2 hours or overnight at 4 °C. After fixation, cells were washed three times with PBS-Tween 20 (0.05% v/v) and blocked with 1 ml PBS-BSA (1%, w/v) in each plate for 30 min incubation at 37°C. Cells were washed three times with PBS-Tween 20. Then first antibody specific to
20 *Helicobacter pylori* raised in rabbits were added (1:600 dilution) and kept for 1 hour at 37°C. Extra unbound first antibody was washed off by washing three times with PBS-Tween20. The secondary antibody which is the FITC

conjugate with mouse monoclonal anti-rabbit IgG (Sigma-Aldrich, Inc) were added (1:100 dilution) and cells were incubated for 1 hour at 37°C. cells were washed well with PBS-Tween 20 at least 3 times. After drying, cells were observed under a microscope (NIKON FDX-35, EX 450-490 nm, DM 5 505, BA 520 nm) and pictures were taken.

FIG. 3 is a photograph showing the inhibition of adherence of *Helicobacter pylori* to gastric cells by PL bacteria, wherein the part showing the yellow color in FIG. 3 is *Helicobacter pylori* labeled with FITC conjugated antibody under a fluorescence microscope. FIG. 3a is the MKN-45 cell, FIG. 3b is the MKN-45 cell treated with *Helicobacter pylori*, FIG. 3c is the MKN-45 cells treated with the surviving PL9001 bacterium and *Helicobacter pylori*, FIG. 3d is the MKN-45 cells treated with the non-viable PL9001 bacteria and *Helicobacter pylori*, FIG. 3e is the MKN-45 cell treated with the surviving PL9003 bacteria and *Helicobacter pylori*, FIG. 3f is the MKN-45 cells treated with the non-viable PL9003 bacteria and *Helicobacter pylori*, FIG. 3g is the MKN-45 cells treated with the surviving PL9004 bacteria and *Helicobacter pylori*, FIG. 3h is the MKN-45 cells treated with the non-viable PL9004 bacteria and *Helicobacter pylori*, FIG. 3i is the MKN-45 cells treated with the surviving PL9005 bacteria and *Helicobacter pylori*, FIG. 3j is the MKN-45 cells treated with the non-viable PL9005 bacteria and *Helicobacter pylori*, FIG. 3k is the MKN-45 cells treated with the surviving PL9006 bacteria and

Helicobacter pylori, and FIG. 3I is the MKN-45 cells treated with the non-viable PL9006 bacterium and *Helicobacter pylori*. In florescence microscopy, MKN-45 cell shows green and *Helicobacter pylori* shows yellow. In a black and white photograph, *Helicobacter pylori* shows more brightly
5 than MKN-45 cells.

It was confirmed that the surviving PL bacteria and non-viable PL bacteria inhibit the adhesion of *Helicobacter pylori* to gastric cells. Therefore, PL bacteria prevent the inhabitation of *Helicobacter pylori* in the stomach so that it can treat and protect from *Helicobacter pylori*-associated
10 disease.

EXAMPLE 3. The adhesion assay conducted with PL bacteria on gastric cell lines

Bacteria having biology activity and non-viable bacteria prepared by boiling at 75°C for 15 min were prepared. The adhesion assay was
15 conducted with the surviving bacteria and the non-viable bacteria on MKN-45 (Human Gastric Adenocarcinoma).

MKN-45 cells were cultured in an RPMI-1640 medium (Gibco-BRL, New York, N.Y.) containing 2 g/L sodium bicarbonate, 10% heat-inactivated fetal bovine serum, and an antibiotic-antimycotic, pH 7.2. A cell monolayer
20 was prepared in a 30 mm dish (Nunc Inc., Roskilde, Denmark) for Gram staining by inoculating 3×10^5 viable cells into a 2 ml culture medium. The medium was replaced every other day. Cells were used for binding at late

post-confluence after 6 days of culture.

After six days the confluent monolayer was washed twice with 2 ml of PBS (phosphate buffered saline) before the adhesion assay. 100 μ L PL bacteria (1×10^7 cfu/ml) were added to 2 ml of a complete medium. This
5 suspension was added to each dish and incubated at 37°C in 5% CO₂-95% air. After incubation for 60 min, the monolayer was washed twice with sterile PBS (pH 7.4), and fixed with methanol. After being stained with Gram staining, cells were examined under a light microscope.

FIG. 4 is photograph showing the adhesion of the surviving PL
10 bacteria and the non-viable PL bacteria to gastric cell, wherein "a" is the surviving PL9001 bacterium, "a'" is the non-viable PL9001 bacterium, "b" is the surviving PL9003 bacterium, "b'" is the non-viable PL9003 bacterium, "c" is the surviving PL9004 bacterium, "c'" is the non-viable PL9004 bacterium, "d" is the surviving PL9005 bacterium, "d'" is the non-viable PL9005
15 bacterium, "e" is the surviving PL9006 bacterium, and "e'" is the non-viable PL9006 bacterium.

Therefore, PL bacteria of the present invention have excellent adherent activity for gastric cells regardless of their biological activity.

EXAMPLE 4

20 PL bacteria were treated with 5 M LiCl₂ (Mark S. TURNER, Peter TIMMS, Louise M. HAFNER, and Philip M. GIFFARD (1997) Journal of Bacteriology, vol. 179, No. 10, p. 3310-3316) to remove the protein of the

cell walls, and the adhesion assay on gastric cells was tested. The number of adhered PL bacteria was measured with a light microscope, and the values are represented in Table 7.

The group treated with LiCl_2 shows a decrease of adherence of bacteria, therefore it is confirmed that protein of cell walls of PL bacteria are involved in adhesion.

(Table 7)

		The observed Bacteria No.	Ratio (%)
The untreated group with LiCl_2		219.5/fields	100
The treated group with LiCl_2	PL9001	10.96/fields	54.8
	PL9003	100.0/fields	50.0
	PL9004	110.0/fields	55.0
	PL9005	98.0/fields	49.0
	PL9006	100.0/fields	50.0

EXAMPLE 5. Test of inhibiting *Helicobacter pylori* growth.

(1) Test of inhibiting *Helicobacter pylori* growth using a culture filtrate of PL bacteria.

The PL bacteria (*Lactobacillus coprophilus* PL 9001(KCCM-10245), *Enterococcus durans* PL 9002(KCCM-10246), *Streptococcus faecalis* PL 9003(KCCM-10247), *Lactobacillus coprophilus* PL 9004(KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251)) was cultured for 24 h at 37°C in MRS broth, and the supernatant was prepared as culture filtrate by centrifugation.

An inhibition test of *Helicobacter pylori* growth using the culture filtrate of PL bacteria was performed.

Wells were formed on a Brucella solid medium using a sterilized Pasteur pipet. *Helicobacter pylori* spread on the medium and the culture filtrate of PL bacteria was added to the well. After incubating in a CO₂ incubator (5% to 10% CO₂) for 2 days, the diameter of the zone that *Helicobacter pylori* growth inhibited due to the inhibitory material of the culture filtrate was measured.

FIG. 5 is a photograph showing that the culture filtrate has growth-inhibitory activity against *Helicobacter pylori*, and FIG. 6 shows a diameter of the growth-inhibited-zone of *Helicobacter pylori* by the culture filtrate. Therefore, it was confirmed that PL bacteria prepared by the EXAMPLE 1 secreted material that inhibited *Helicobacter pylori* growth.

(2) Test of inhibiting *Helicobacter pylori* growth by lyophilized culture filtrate of PL bacteria.

4 ml of MRS culture filtrate of PL bacteria was mixed with 2 ml of distilled MQ and the mixture was lyophilized. The lyophilized culture filtrate was suspended on 6 ml of skim milk (10 mg/ml) and after 1 ml of the solution was inoculated with 10 ul of *Helicobacter pylori* (OD₆₂₅=1.0), it was incubated for 1hr at 37°C in a CO₂ incubator. The incubated solution was centrifuged (5000 rpm, 10min) and the supernatant was diluted by half with distilled MQ. The quantity of urease of the diluted solution was measured by an indolphenol method which assay the amount of ammonia produced from urea by urease. Since the quantity of urease is proportional to the amount

of *Helicobacter pylori*, the quantity of ammonia is proportional to the number of *Helicobacter pylori* in the incubated solution. Through these methods, growth inhibition of *Helicobacter pylori* by culture filtrate of PL bacteria was analyzed.

5 FIG. 7 is a graph showing the growth-inhibition degree of *Helicobacter pylori* by culture filtrate of PL bacteria, and it shows that the lyophilized culture filtrate of PL bacteria maintains the growth inhibition activity for *Helicobacter pylori*.

EXAMPLE 6. Toxicity test

10 The oral toxic test for the PL bacteria of EXAMPLE 1 was based on the "Toxic Test Standard for Drugs (1999. 12. 22)," Notification No. 1999-61 of the Korean Food and Drug Administration.

Tests were conducted on Sprague-Dawley rats, which were 5 weeks old (female : 100 ~ 120g / male : 110 ~ 130g). The rats were reared in
15 260 × 420 × 180 mm (W × L × H) cages at 23 ± 2°C and 50 ± 10 humidity.

Referring to established rule No. 10 of the Korean Food and Drug Administration, when the value of LD₅₀ is up to 5,000 mg/kg in an oral test, the material is a low toxicity material in a body. Therefore, in the
20 experiment, to calculate the value of LD₅₀ of PL bacteria, the dosage was set to 5,000 mg/kg (20 ml/kg B.W.) 5,000 mg/kg was the maximum dosage given to the rats to determine the toxicity of the material.

(Table 8)

Bacteria	Sex	number	Dosage(mg/kg B.W)	Dosage(ml/kg B.W)
I(PL 9001)	Male	5	5,000	20
	Female	5		
II(PL 9002)	Male	5	5,000	20
	Female	5		
III(PL 9003)	Male	5	5,000	20
	Female	5		
IV(PL 9004)	Male	5	2,500	20
	Female	5		
V(PL 9004)	Male	5	5,000	20
	Female	5		
VI(control)	Male	5	0	20
	Female	5		

After 14 days, no deaths were observed. Thus the value of LD₅₀ could not be estimated.

Table 9 shows the weight change of the test group, and there were
5 no significant changes.

(Table 9)

Sex	Days	I	II	III	IV	V	VI
M	0	120.1± 3.10	120.2± 3.08	120.1± 3.08	120.4± 2.86	120.2± 3.25	120.1± 3.31
	3	142.6± 6.81	146.9± 4.15	140.5± 5.46	140.7± 8.85	145.4± 7.73	143.0± 3.68
	7	162.7± 7.96	161.0± 5.87	155.5± 2.92	156.6± 11.59	162.1± 5.64	159.8± 3.06
	14	212.3± 8.40	193.1± 18.03	194.9± 9.14	201.1± 22.72	195.5± 11.14	193.6± 16.24
F	0	114.0± 4.34	113.9± 4.11	113.9± 4.17	113.5± 4.41	113.5± 4.45	113.6± 4.50
	3	134.3± 6.81	132.6± 4.15	131.0± 5.46	132.5± 8.85	129.8± 7.73	129.2± 3.68
	7	155.4± 8.01	151.5± 5.69	149.5± 5.86	152.2± 7.55	148.6± 5.81	148.8± 6.57
	14	199.8± 8.16	192.7± 10.87	192.5± 10.0	193.8± 12.06	194.7± 7.55	192.6± 9.35

In addition, all the rats were examined to determine abnormalities,

but there was no symptoms visible to ordinary sight (Table 10).

(Table 10)

Organ		I	II	III	IV	V	VI
Brain	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Kidney-Left	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Kidney-Right	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Heart	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Lung	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Spleen	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Liver	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Stomach	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Intestine	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Pancreas	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Adrenal gland(left)	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Adrenal gland(right)	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Pituitary gland	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Ovary-L	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Ovary-R	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5
Other organs	No. of observations	5	5	5	5	5	5
	No gross findings	5	5	5	5	5	5

As shown above, PL bacteria of the EXAMPLE is safe without oral toxicity and side effects.

5 EXAMPLE 7. Growth inhibition test for bacteria that cause food poisoning

An MRS culture filtrate of each PL bacterium was mixed with a 2 X concentrated BHI medium. Each of *Salmonella typhimurium*, *Salmonella*

Enteriditis, *Escherichia coli* O157:H7, *Aeromonas hydrophila*, and *Listeria monocytogenes* cultured in BHI (Brain Heart Infusion: Brain Heart, Infusion form, 6.0 g, Peptic digest of animal tissue 6.0 g, sodium chloride 5.0 g, dextrose 3.0 g, pancreatic digest of gelatin 14.5 g, disodium phosphate 2.5 g) were inoculated with a 1% final concentration of the mixture and incubated at 37°C. After 5 hr and 12 hr, the number of surviving bacteria was measured. For measurement, *Listeria monocytogenes* (L.M) was incubated in a blood-agar plate, and *Salmonella typhimurium* (S.T), *Salmonella Enteriditis* (S.E), *E. coli* O157:H7 (O157), and *Aeromonas hydrophila* (A.H) were incubated in a MacConkey medium, and the number of surviving bacteria was measured at O.D₆₀₀.

Table 11 shows an O.D₆₀₀ of bacteria that cause food poison by a PL 9001 culture filtrate.

(Table 11)

Condition	0 hour	5 hour	24 hour
L.M	0.055	0.487	0.335
L.M + culture filtrate (PL9001)	0.059	0.069	0.068
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL9001)	0.063	0.073	0.072
S.E	0.053	0.572	0.966
S.E+ culture filtrate (PL9001)	0.059	0.073	0.07
O157	0.051	0.408	0.468
O157+ culture filtrate (PL9001)	0.06	0.074	0.065
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL9001)	0.069	0.093	0.083

15

FIG. 8 is a graph showing growth inhibition of bacteria that cause food poisoning by *Lactobacillus coprophilus* PL 9001, and it shows the

results of Table 11 (absorbance of $O.D_{600}$ at 0 hr was calculated to 100). As shown above in Table 11 and in FIG. 8, *Lactobacillus coprophilus* PL 9001 produced materials that inhibit the growth of bacteria that cause food poisoning.

5 Also, *Enterococcus durans* PL 9002 inhibited the growth of bacteria that cause food poisoning. Inhibition results are shown in Table 12 and FIG.

9.

(Table 12)

Condition	0 hour	5 hour	24 hour
L.M	0.055	0.487	0.335
L.M+ culture filtrate (PL 9002)	0.059	0.071	0.17
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL 9002)	0.064	0.073	0.196
S.E	0.053	0.572	0.966
S.E+ culture filtrate (PL 9002)	0.061	0.074	0.167
O157	0.051	0.408	0.468
O157+ culture filtrate (PL 9002)	0.59	0.073	0.17
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL 9002)	0.065	0.086	0.235

The inhibition effect of *Streptococcus faecalis* PL 9003 for growth of
 10 bacteria that cause food poisoning was measured. Table 13 shows $O.D_{600}$ of bacteria that cause food poisoning in a culture solution containing *Streptococcus faecalis* PL 9003, and the data of Table 13 are shown in the graph of FIG 10, showing that *Streptococcus faecalis* PL 9003 produced materials that inhibit the growth of bacteria that cause food poisoning.

15

(Table 13)

Condition	0 hour	5 hour	24 hour
-----------	--------	--------	---------

L.M	0.055	0.487	0.335
L.M+ culture filtrate (PL 9003)	0.058	0.007	0.075
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL 9003)	0.063	0.074	0.084
S.E	0.053	0.572	0.966
S.E+ culture filtrate (PL 9003)	0.062	0.074	0.082
O157	0.051	0.408	0.468
O157+ culture filtrate (PL 9003)	0.058	0.07	0.071
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL 9003)	0.066	0.085	0.094

The inhibition effect of *Lactobacillus coprophilus* PL 9004 for the growth of bacteria that cause food poisoning was measured. Table 14 shows O.D₆₀₀ of bacteria that cause food poisoning in a culture solution containing *Lactobacillus coprophilus* PL 9004, and the data of Table 14 are shown in the graph of FIG. 11 of data of Table 14, showing that *Lactobacillus coprophilus* PL 9004 produced materials that greatly inhibit the growth of bacteria that cause food poisoning, but the inhibition effect was decreased.

(Table 14)

Condition	0 hour	5 hour	24 hour
L.M	0.055	0.487	0.335
L.M+ culture filtrate (PL 9004)	0.06	0.075	0.287
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL 9004)	0.063	0.071	0.266
S.E	0.053	0.572	0.966
S.E+ culture filtrate (PL 9004)	0.061	0.073	0.227
O157	0.051	0.408	0.468
O157+ culture filtrate (PL 9004)	0.062	0.073	0.267
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL 9004)	0.065	0.087	0.251

The inhibition effect of *Lactobacillus fermentum* PL 9005 for the growth of bacteria that cause food poisoning was measured. Table 15

below shows $O.D_{600}$ of bacteria that cause food poisoning in a culture solution containing *Lactobacillus fermentum* PL 9005, and FIG 12 is a graph showing the growth inhibition for Listeria. *Lactobacillus fermentum* PL 9005 produced materials that greatly inhibit the growth of bacteria that cause food poisoning.

(Table 15)

Condition	0 hr	5 hr	24 hr
L.M	0.055	0.487	0.335
L.M+ culture filtrate (PL 9005)	0.06	0.068	0.413
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL 9005)	0.065	0.074	0.434
S.E	0.053	0.572	0.966
S.E+ culture filtrate (PL 9005)	0.062	0.075	0.448
O157	0.051	0.408	0.468
O157+ culture filtrate (PL 9005)	0.06	0.077	0.389
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL 9005)	0.067	0.087	0.245

The inhibition effect of *Lactobacillus fermentum* PL 9006 for the growth of bacteria that cause food poisoning was measured. Table 16 shows $O.D_{600}$ of bacteria that cause food poisoning in a culture solution containing *Lactobacillus fermentum* PL 9006, and FIG 13 is a graph showing the growth inhibition for Listeria. *Lactobacillus fermentum* PL 9006 produced materials that greatly inhibit the growth of bacteria that cause food poisoning.

(Table 16)

Condition	0 hr	5 hr	24 hr
L.M	0.055	0.487	0.335
L.M+ culture filtrate (PL 9006)	0.058	0.068	0.283
S.T	0.053	0.538	0.748
S.T+ culture filtrate (PL 9006)	0.061	0.073	0.303
S.E	0.053	0.572	0.966

S.E+ culture filtrate (PL 9006)	0.065	0.069	0.428
O157	0.051	0.408	0.468
O157+ culture filtrate (PL 9006)	0.063	0.077	0.435
A.H	0.077	0.418	0.432
A.H+ culture filtrate (PL 9006)	0.067	0.087	0.112

As mentioned above, PL bacteria (*Lactobacillus coprophilus* PL 9001, *Enterococcus durans* PL 9002, *Streptococcus faecalis* PL 9003, *Lactobacillus coprophilus* PL 9004, *Lactobacillus fermentum* PL 9005 and *Lactobacillus fermentum* PL 9006) of the present invention produced materials that inhibit the growth of bacteria that cause food poisoning. Also, the decreased inhibition effect is caused by depleting material derived from culture filtrate, and then it can maintain the growth inhibition effect for bacteria that cause food poisoning by using PL bacteria.

EXAMPLE 8. Growth inhibition effect for bacteria that cause acne

Propionibacterium acne is a normal bacteria that exists on skin and causes acne. *Propionibacterium acne* (KCTC 3314) was inoculated into 5 ml of actinomyces broth, and after covering by paraffin-oil, *Propionibacterium acne* was anaerobically incubated for 2 days (BR Vowels, S Yang, JJ Leyden. 1995. Induction of proinflammatory cytokines by a soluble factor of *Propionibacterium acnes*: Implications for chronic inflammatory acne. Infection and Immunity 63: 3158-3165). 45 ml of PL culture filtrate bred on a MRS liquid medium was mixed with 5 ml of *Propionibacterium acnes* culture solution. 50 ul of the mixture were inoculated into 5 ml of a fresh medium, an actinomyces broth, and cultured anaerobically for 2 days without shaking.

For measurement of survival *Propionibacterium acnes*, the culture solution

was diluted (When $OD_{600}=2.4$, 9.6×10^8 CFU/ml) with a diluent containing 0.05% L-cysteine for anaerobic bacteria, and 100 ul of the diluted solution was inoculated onto an actinomyces agar plate. After culture in an anaerobic jar at 37°C for 6-7 days, the colony on the plate was measured as
5 the number of survival *Profionibacterium acnes*.

FIG. 14 is a graph showing the growth inhibition activity for bacteria that cause acne by PL bacteria, and *Lactobacillus coprophilus* PL 9001 did not inhibit nearly the growth of *Profionibacterium acnes*; *Enterococcus durans* PL 9002 and *Streptococcus faecalis* PL 9003 completely inhibited the
10 growth of *Profionibacterium acnes*; and *Lactobacillus coprophilus* PL 9004, *Lactobacillus fermentum* PL 9005 and *Lactobacillus fermentum* PL 9006 inhibited the growth of *Profionibacterium acnes*.

EXAMPLE 9. Growth inhibition test for anaerobic bacteria.

Clostridium perfringens was inoculated into BHI liquid broth containing
15 Hemin (0.01 g/L) and L-cysteine (0.5 g/L) and it was anaerobically cultured at 37°C for 24 hours (Balows A Hausler WJ Herrmann KL Isenberg HD Shadomy HJ. Chapter 50. Clostridium. p505-521. Manual of Clinical Microbiology, 5th ed. ASM Washington D.C. U.S.A.). A culture filtrate of PL bacteria was mixed with the same volume of 2X concentrated BHI broth, and
20 a 1% final concentration of *Clostridium perfringens* was inoculated into the mixture. After covering the mixture with parafilm-oil, it was anaerobically cultured at 37°C. After 24 hours, the culture solution was diluted and

inoculated onto a blood agar plate. The plate was incubated for 37 days and 48 days, in an anaerobic incubator. The survival rate of *Clostridium perfringens* was measured and is presented by log.

(Table 17)

	Number of survival bacteria (log cfu)	
	0 hrs	24 hrs
<i>Clostridium perfringens</i>	6.60	8.30
<i>Lactobacillus coprophilus</i> PL 9001	6.60	7.60
<i>Enterococcus durans</i> PL 9002	6.78	7.66
<i>Streptococcus faecalis</i> PL 9003	6.60	7.48
<i>Lactobacillus coprophilus</i> PL 9004	6.90	8.00
<i>Lactobacillus fermentum</i> PL 9005	6.60	7.78
<i>Lactobacillus fermentum</i> PL 9006	6.60	7.41

5 It was confirmed that all the PL bacteria produced materials that inhibit growth of anaerobic bacteria.

EXAMPLE 10. Improving effect of immunity

The immune-improving effect of PL bacteria was tested. When PL bacteria was added to a macrophage, the amount of the producing cytokine
 10 (TNF- α , IL-6) was quantified.

50 ul of a coating buffer containing purified rabbit anti-TNF- α or rabbit anti-IL-6 was coated within a microplate well and the microplate was maintained at -4°C for 12 hr. The microplate well was washed three times with a PBST (tween80+phosphate buffer) solution and reacted with 3% BSA-
 15 PBST, for 30 min. The microplate well was washed four times with PBST and reacted with 50 ul of TNF- α -standard/IL-6 standard (recombinant mouse TNF- α /IL-6, SEROTEC, UK) or macrophage supernatant cultured

in a medium containing PL bacteria, at 37°C for 1 hr. The microplate well was washed four times with PBST, once with distilled water, and it was reacted with 50 ul of a mixture prepared by adding 3% BSA-PBST to biotinylated rat anti-mouse TNF- α or biotinylated rat anti-mouse TNF- α /IL-6 (biotinylated rat anti-mouse TNF- α /IL-6, 1 ug/ml), at room temperature for 1 hr. The microplate well was washed six times with PBST, once with distilled water, and reacted with streptavidin peroxidase for one hour. The microplate well was washed eight times with PBST, twice with distilled water, and reacted with TMB substrate (25 ml citric-phosphate buffer+400 ul TMB stock+100 ul of 1% H₂O₂) at room temperature until color showed. The reaction was stopped to add 6 N of H₂SO₄, and for measurement of immune reactivity, adsorption at OD₄₅₀ was observed.

FIG. 15 and Table 18 show an immune-improving effect of PL bacteria, and a PL bacteria increase of the production of TNF- α and IL-6.

15

(Table 18)

		Adsorption at O.D. ₄₅₀		Increase ratio %
		Control	Test	
PL9001	TNF- α	15	136	9.06
	IL-6	0.5	21.5	43
PL9003	TNF- α	15	58	3.87
	IL-6	0.5	16.5	33
PL9004	TNF- α	15	88	5.87
	IL-6	0.5	18	36
PL9005	TNF- α	15	38	2.3
	IL-6	0.5	21.5	43
PL9006	TNF- α	15	59	3.93
	IL-6	0.5	21.4	42.8

Therefore, PL bacteria of the present invention can be applied to

promote immunity, and more particularly, it can be used for health food and as a treatment drug that promotes the health of aged persons and children.

EXAMPLE 11. Test for acid-resistance and bile-resistance

An MRS medium containing cysteine was optimized according to pH (7, 4.5 or 4.0) with 4 N HCl and 0.1 N NaOH, and was sterilized. To observe an effect of bile, oxgall powder was added to the medium in 0%, 0.25%, 0.50% concentrations and it was sterilized. Each PL bacteria was inoculated into the medium with a total of 1% concentration, and optical density was determined at 620 nm at 24-hours intervals. Table 19 shows the acid-resistance and bile-resistance. It was shown that PL 9003 was weak in acid, but PL 9001, PL 9002, PL 9004, PL 9005, and PL 9006 had strong acid-resistant activity, and all the PL bacteria had strong bile-resistant activity. Therefore, all PL bacteria are safe in the stomach and intestine (Conway PL Gorback SL goldin BR. 1987. Survival of lactic acid bacteria in the human stomach and adhesion to intestinal cells. J. Dairy Sci. 70: 1-12. : Ibrahim SA Bezkorovainy A. 1993. Survival of bifidobacteria in the presence of bile salt. J. Sci. Food Agric. 62: 351-354).

(Table 19)

	Time	PL 9001	PL 9002	PL 9003	PL 9004	PL 9005	PL 9006
pH 7	24 hr	1.40	1.20	1.00	1.40	1.50	1.50
	48 hr	1.50	1.50	1.20	1.50	1.50	1.50
pH 5.0	24 hr	1.00	0.70	0.31	0.90	1.50	1.50
	48 hr	1.50	0.80	0.45	1.10	1.50	1.50
pH 4.5	24 hr	0.18	0.21	0.06	0.18	1.50	1.50
	48 hr	0.70	0.28	0.09	0.15	1.50	1.50
Oxgall 0.25%	24 hr	1.40	1.20	1.00	1.20	1.10	1.20
	48 hr	1.50	1.50	1.40	1.50	1.40	1.20

Oxgall 0.50%	24 hr	1.10	1.20	1.10	0.85	0.75	0.70
	48 hr	1.40	1.50	1.40	1.00	0.76	0.70

EXAMPLE 12. Antibiotic-resistance

Antibiotic-resistance for PL bacteria prepared by the EXAMPLE was observed. PL bacteria were inoculated into an MRS solid medium and a filter (diameter 6 mm) containing the antibiotic of Table 20 was put on the medium. After incubation for 24–48 hr, the diameter of the growth inhibited-zone formed by antibiotics was determined. A decreased diameter of the growth inhibited-zone means that PL bacteria have a resistance to the antibiotics. The results are presented in Table 20.

10 (Table 20)

Antibiotics	PL9001	PL9002	PL9003	PL9004	PL9005	PL9006
Penicillin (10 IU/E/UI)	20 mm	20 mm	24 mm	23 mm	30 mm	30 mm
Ampicillin (10 ug)	30 mm	24 mm	26 mm	30 mm	30 mm	32 mm
Cephalothin (30 ug)	30 mm	20 mm	20 mm	24 mm	30 mm	30 mm
Gentamycin (10 ug)	13 mm	-	-	12 mm	15 mm	16 mm
Vancomycin (30 ug)	-	20 mm	16 mm	-	-	-
Erythromycin (15 ug)	21 mm	-	-	22 mm	26 mm	29 mm
Tetracycline (30 ug)	26 mm	30 mm	10 mm	24 mm	27 mm	30 mm

As mentioned above, PL bacteria have high antibiotic resistance.

EXAMPLE 13. The adhesion assay conducted with PL bacteria on intestinal cell lines

Caco-2 cells (intestinal cell line) were cultured in an RPMI-1640 medium (Gibco-BRL, New York, N.Y.) containing 2 g/L sodium bicarbonate, 10% heat-inactivated fetal bovine serum, and antibiotic-antimycotic, pH 7.2. A cell monolayer was prepared in a 30 mm dish (Nunc Inc., Roskilde,

Denmark) for Gram staining by inoculating 3×10^5 viable cells into a 2 ml culture medium. The medium was replaced every other day. Cells were used for binding at late post-confluence after 6 days of culture.

After six days the confluent monolayer was washed twice with 2 ml
 5 PBS (phosphate buffered saline) before the adhesion assay. 100 μ L PL bacteria (1×10^7 cfu/ml) were added to 2 ml of a complete medium. This suspension was added to each dish and incubated at 37°C in 5% CO₂-95% air. After incubation for 60 min, the monolayer was washed twice with sterile PBS (pH 7.4), and fixed with methanol. After being stained with Gram
 10 staining, cells were examined under a light microscope.

The number of observed bacteria within 20 fields was measured, and when their number was over 400 it was determined that adhesion occurred (C.N. JACOBSEN, V. ROSENFELDT, NIELSEN, A.E. HAYFORD, P.L. MØLLER, K.F. MICHAELSEN, A. PAERREGAARD, B. SANDSTRÖM, M.
 15 TVEDE, and M. JAKOBSEN (1999) *Applied and Environmental Microbiology*, p.4949-4956 ; R. Bibiloni, P.F. Pérez, and G.L. DeAntoni (1999) *Anaerobe* 5, 483-485).

Table 21 shows a number of adhered PL bacteria. PL bacteria of the present invention are shown to have an adherence activity on intestinal
 20 cells.

(Table 21)

Name of the added PL bacteria	Attached bacteria No. on Caco-2
<i>Bifidobacterium infantis</i>	>800/20 fields

<i>Bifidobacterium longum</i>	>800/20 fields
PL9001	>800/20 fields
PL9003	>800/20 fields
PL9004	>800/20 fields
PL9005	>800/20 fields
PL9006	>800/20 fields

FIG. 16 is photograph showing the adhered PL bacteria on CaCo-2 cell lines. The adherence of PL9001 (b), PL9003 (c), PL9004 (d), PL9005 (e) and PL9006 (f) on intestinal cell lines was observed and compared to normal intestinal cell lines (a).

Bifidobacterium infantis and *Bifidobacterium longum* are the known adherent bacteria on intestinal cell lines, and compared to them, PL bacteria of the present invention are excellent adherent bacteria on intestinal cells and can therefore be applied to clean intestines.

10

EXAMPLE 16. Preparation of cosmetics

Preparation of toner

0.02 wt% powder prepared by drying PL 9001, 5.0 wt% glycerin, 3.0 wt% 1,3-butylene glycol, 1.0 wt% PEG1500, 0.1 wt% allantoin, 0.3 wt% DL-pantenol, 0.02 wt% EDTA-2NA, 0.04 wt% benzopenon-9, 5.0 wt% sodium hyaluronate, 10.0 wt% ethanol, 16 wt% octyldodecene, 20 wt% polysorbate, antiseptic, pigment, and distilled water were mixed and toner was prepared.

Preparation of lotion

0.01 wt% of powder prepared by drying PL 9001, 5.0 wt% glycerin, 3.0 wt% 1,3-butyleneglycol, 5.0 wt% sodium hyaluronate, 10.0 wt% ethanol, 60wt% polysorbate, 1.5 wt% glycerylstearate 1.5 wt% stearyl alcohol, 1.5

20

wt% lanolin, 0.5 wt% sorbitan stearate, 1.0 wt% vegetable oils, 5.0 wt% mineral oil, 5.0 wt% squalene, 2.0wt% trioctanoin, 0.8wt% dimethicone, 0.5 wt% tocopherol acetate, 0.12 wt% carboxyvinylpolymer, 0.12 wt% triethanolamine, antiseptic, pigment, perfume, and distilled water were mixed
5 and lotion was prepared.

Applicant's or agent's file reference	OPP010966KR	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>8</u> , line <u>1</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <p style="text-align: center;">KOREA CULTURE CENTER OF MICROORGANISMS</p>	
Address of depositary institution (including postal code and country) <p style="text-align: center;">361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL, 120-091, Republic of Korea</p>	
Date of deposit <p style="text-align: center;">December 2, 2000</p>	Accession Number <p style="text-align: center;">KCCM-10245</p>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer <p style="text-align: center;">[Signature]</p>	Authorized officer

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To, Yeon-hee Lee

Department of Biology and Culture Collection
of Antibiotic Resistant Microbes, College of
Natural Science, Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR :

Lactobacillus coprophilus PL 9-1

Accession number given by the
INTERNATIONAL DEPOSITARY AUTHORITY :

KCCM - 10245

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☐ a scientific description

☐ a proposed taxonomic designation

(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

the microorganism identified under I above was received by this international Depositary Authority on Dec. 2, 2000. and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on Jan. 31, 2001.

IV. INTERNATIONAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms

Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s) :

Date: Jan. 31, 2001



BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yeonhee lee
Department of Biology,
Seoul woman's University,
Seoul 139-774,
Korea


The enclosed communication has been received by this International Depositary Authority
on Jun. 21, 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Sondaemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s) :

Date: Jun. 25, 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION


pursuant to Rule 8.1

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul, 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10245	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> ¹ Scientific description:	
<input type="checkbox"/> ² Last preceding scientific description (if any):	
<input type="checkbox"/> ² Proposed taxonomic designation:	
	Lactobacillus PL 9001
<input type="checkbox"/> ² Last preceding proposed taxonomic designation (if any):	
	Lactobacillus CCARM 9-1

¹. Mark with a cross if additional information is given on an attached sheet.

². Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
The undersigned 111	
<input type="checkbox"/> requests	
<input type="checkbox"/> does not request	
the attestation referred to in Rule 8.2	
IV. DEPOSITOR	
Name: Yeonhee lee	Signature ⁴ : 
Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea	Date: June. 21. 2001

³. Mark with a cross the application box.

⁴. Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

Applicant's or agent's file reference	OPP010966KR	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>8</u> , line <u>2</u>	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution KOREA CULTURE CENTER OF MICROORGANISMS	
Address of depositary institution (including postal code and country) 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL, 120-091, Republic of Korea	
Date of deposit December 2, 2000	Accession Number KCCM-10246
C. ADDITIONAL INDICATIONS (leave blank if not applicable)	
This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer V. J. H. S.	Authorized officer

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To, Yeon-hee Lee

Department of Biology and Culture Collection
of Antibiotic Resistant Microbes, College of
Natural Science, Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR :

Lactobacillus plantarum PL 9-2

Accession number given by the
INTERNATIONAL DEPOSITARY AUTHORITY :

KCCM - 10246

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☐ a scientific description

☐ a proposed taxonomic designation

(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

the microorganism identified under I above was received by this international Depositary
Authority on Dec. 2, 2000. and a request to convert the original deposit to a deposit
under the Budapest Treaty was received by it on Jan. 31, 2001.

IV. INTERNATIONAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms

Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s) :

Date: Jan. 31, 2001



BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yeonhee lee
Department of Biology,
Seoul woman's University,
Seoul 139-774,
Korea

The enclosed communication has been received by this International Depositary Authority
on Jun. 21. 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaeemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s)

Date: Jun. 25. 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE


COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION
pursuant to Rule 2.1

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10246	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> Scientific description:	
<input type="checkbox"/> Last preceding scientific description (if any):	
<input type="checkbox"/> Proposed taxonomic designation: Enterococcus PL 9002	
<input type="checkbox"/> Last preceding proposed taxonomic designation (if any): Lactobacillus CCARM 9-2	

¹ Mark with a cross if additional information is given on an attached sheet.

² Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
<p>The undersigned 111</p> <p><input type="checkbox"/> requests</p> <p><input type="checkbox"/> does not request</p> <p>the attestation referred to in Rule 8.2</p>	
IV. DEPOSITOR	
<p>Name: Yeonhee lee</p> <p>Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea</p>	<p>Signature¹ : </p> <p>Date: Jun. 21. 2001</p>

¹ Mark with a cross the application box.

² Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To, Yeon-hee Lee

Department of Biology and Culture Collection
of Antibiotic Resistant Microbes, College of
Natural Science, Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR :

Lactobacillus para.paracasei 1 PL 9-3

Accession number given by the
INTERNATIONAL DEPOSITARY AUTHORITY :

KCCM - 10247

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☐ a scientific description

☐ a proposed taxonomic designation

(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

the microorganism identified under I above was received by this international Depositary Authority on Dec. 2, 2000. and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on Jan. 31, 2001.

IV. INTERNATIONAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms

Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s) :

Date: Jan. 31, 2001



BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yeonhee lee
Department of Biology,
Seoul woman's University,
Seoul 139-774,
Korea

The enclosed communication has been received by this International Depositary Authority
on Jun. 21. 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s)

Date: Jun. 25. 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION


pursuant to Rule 2.1.

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10247	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> Scientific description:	
<input type="checkbox"/> Last preceding scientific description (if any):	
<input type="checkbox"/> Proposed taxonomic designation: Enterococcus PL 9003	
<input type="checkbox"/> Last preceding proposed taxonomic designation (if any): Lactobacillus CCARM 9-3	

¹ Mark with a cross if additional information is given on an attached sheet.

² Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
The undersigned <u>111</u>	
<input type="checkbox"/> requests	
<input type="checkbox"/> does not request	
the attestation referred to in Rule 8.2	
IV. DEPOSITOR	
Name: Yeonhee lee	Signature: 
Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea	Date: June. 21. 2001

* Mark with a cross the application box.

* Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

Applicant's or agent's file reference	OPP010966KR	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>8</u> , line <u>3</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <p style="text-align: center;">KOREA CULTURE CENTER OF MICROORGANISMS</p>	
Address of depositary institution (including postal code and country) <p style="text-align: center;">361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL, 120-091, Republic of Korea</p>	
Date of deposit <p style="text-align: center;">December 2, 2000</p>	Accession Number <p style="text-align: center;">KCCM-10248</p>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer <p style="text-align: center;">v/3 in 56</p>	Authorized officer

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To. Yeon-hee Lee

Department of Biology and Culture Collection
of Antibiotic Resistant Microbes, College of
Natural Science, Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Lactobacillus caprophilus</i> PL 9-4	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY : KCCM - 10248
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
the microorganism identified under I above was received by this international Depositary Authority on Dec. 2. 2000. and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on Jan. 31. 2001.	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s) : Date: Jan. 31. 2001

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yoonhee lee
Department of Biology,
Seoul woman's University,
Seoul 139-774,
Korea


The enclosed communication has been received by this International Depositary Authority
on Jun. 21. 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s) :

Date: Jun. 25. 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE


COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION
pursuant to Rule 8.1

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul, 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10248	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> Scientific description:	
<input type="checkbox"/> Last preceding scientific description (if any):	
<input type="checkbox"/> Proposed taxonomic designation: Lactobacillus PL 9004	
<input type="checkbox"/> Last preceding proposed taxonomic designation (if any): Lactobacillus CCARM 9-4	

¹ Mark with a cross if additional information is given on an attached sheet.

² Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
The undersigned <u>111</u>	
<input type="checkbox"/> requests	
<input type="checkbox"/> does not request	
the attestation referred to in Rule 3.2	
IV. DEPOSITOR	
Name: Yeouhee lee	Signature: 
Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea	Date: June. 21. 2001

². Mark with a cross the application box.

⁴. Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

Applicant's or agent's file reference	OPP010966KR	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>8</u> , line <u>4</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <p style="text-align: center;">KOREA CULTURE CENTER OF MICROORGANISMS</p>	
Address of depositary institution (including postal code and country) <p style="text-align: center;">361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL, 120-091, Republic of Korea</p>	
Date of deposit <p style="text-align: center;">March 2, 2001</p>	Accession Number <p style="text-align: center;">KCCM-10250</p>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer <p style="text-align: center;">123 22 46</p>	Authorized officer


BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To. Yeonhee Lee

Department of Biology,
Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Lactobacillus</i> PL 9-5	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10250
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 2, 2001. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority of of authorized Date: Mar. 8, 2001. <div style="text-align: right; margin-top: 20px;">  </div>

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yeonhee Lee
Department of Biology,
Soul woman's University,
Soul 139-774,
Korea

The enclosed communication has been received by this International Depositary Authority
on Jun. 21. 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s) :

Date: Jun. 25. 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE


COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION
pursuant to Rule 8.1

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul, 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10250	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> Scientific description:	
<input type="checkbox"/> Last preceding scientific description (if any):	
<input type="checkbox"/> Proposed taxonomic designation: Lactobacillus PL 9005	
<input type="checkbox"/> Last preceding proposed taxonomic designation (if any): Lactobacillus PL 9-5	

¹ Mark with a cross if additional information is given on an attached sheet.

² Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
<p>The undersigned ¹¹¹</p> <p><input type="checkbox"/> ³ requests</p> <p><input type="checkbox"/> ³ does not request</p> <p>the attestation referred to in Rule 8.2</p>	
IV. DEPOSITOR	
<p>Name: Yoonhee lee</p> <p>Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea</p>	<p>Signature⁴ : </p> <p>Date: June. 21. 2001</p>

³ Mark with a cross the application box.

⁴ Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

Applicant's or agent's file reference OPP010966KR	International application No.
---	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>8</u> , line <u>5</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution KOREA CULTURE CENTER OF MICROORGANISMS	
Address of depositary institution (including postal code and country) 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL, 120-091, Republic of Korea	
Date of deposit March, 2, 2001	Accession Number KCCM-10251
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<div>For receiving Office use only</div> <div><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div>Authorized officer VJ 2 52</div>	<div>For International Bureau use only</div> <div><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div>Authorized officer</div>


**BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE**

INTERNATIONAL FORM

To: Yeonhee Lee

Department of Biology,
Seoul Woman's University,
Seoul 139-774, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Lactobacillus</i> PL 9-6	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10251
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 2, 2001. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D. Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official Date: Mar. 8, 2001. <div style="text-align: right; margin-top: 10px;">  </div>

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

ATTESTATION CONCERNING THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION AND/OR
PROPOSED TAXONOMIC DESIGNATION

pursuant to Rule 8.2

TO Yeonhee lee
Department of Biology,
Seoul woman's University,
Seoul 139-774,
Korea

The enclosed communication has been received by this International Depositary Authority
on Jun. 21. 2001.

INDUSTRIAL DEPOSITARY AUTHORITY

Name : Korean Culture Center of Microorganisms
Address : 361-221, Yurim B/D
Hongje-1-dong,
Seodaemun-gu,
SEOUL 120-091
Republic of Korea

Signature(s) of person(s) having the power to
represent the International Depositary Authority
or authorized official(s) :

Date: Jun. 25. 2001.



Enclosure: Communication of the later indication or an amendment of the scientific description proposed taxonomic designation pursuant
to Rule 8.1

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE


COMMUNICATION OF THE LATER INDICATION OR AN
AMENDMENT OF THE SCIENTIFIC DESCRIPTION
AND/OR PROPOSED TAXONOMIC DESIGNATION
pursuant to Rule 8.1

TO: KCCM
361-221 Yurim B/D
Hongje-1-dong Seodaemun-gu
Seoul. 120-091
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM 10251	
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION <input type="checkbox"/>	
<input type="checkbox"/> Scientific description:	
<input type="checkbox"/> Last preceding scientific description (if any):	
<input type="checkbox"/> Proposed taxonomic designation: Lactobacillus PL 9006	
<input type="checkbox"/> Last preceding proposed taxonomic designation (if any): Lactobacillus PL 9-6	

¹ Mark with a cross if additional information is given on an attached sheet.

² Mark with a cross the applicable box or boxes.

III. REQUEST FOR ATTESTATION	
The undersigned	
<input type="checkbox"/> requests	
<input type="checkbox"/> does not request	
the attestation referred to in Rule 8.2	
IV. DEPOSITOR	
Name: Yconhee lee	Signature: 
Address: Department of Biology, Seoul women's University, Seoul 139-774, Korea	Date: June. 21. 2001

³. Mark with a cross the application box.

⁴. Where the signature is required on behalf of a legal entity, the typewritten name(s) of the natural person(s) signing on the legal entity should accompany the signature(s).

WHAT IS CLAIMED IS:

1. A bacterium for inhibiting the adhesion of *Helicobacter pylori* to the gastric mucosa, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*,
5 *Streptococcus faecalis*, and *Lactobacillus fermentum*.
2. The bacterium of claim 1, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus* PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus*
10 PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251).
3. The bacterium of claim 1, wherein bacterium is live, dehydrated, or non-viable bacterium.
4. A bacterium for inhibiting the growth of *Helicobacter pylori*,
15 wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.
5. The bacterium of claim 4, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus coprophilus* PL 9001
20 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus* PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250),

and *Lactobacillus fermentum* PL 9006 (KCCM-10251).

6. A composition for inhibiting the growth of bacteria, wherein the composition comprises at least one bacterium selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus*
5 *faecalis*, and *Lactobacillus fermentum*.

7. The composition of claim 6, wherein the composition comprises at least one selected from the group consisting of *Lactobacillus coprophilus* PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus*
10 *coprophilus* PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251).

8. The composition of claim 6, wherein the bacteria is *Helicobacter pylori*, a bacteria that causes food poisoning, a bacteria that causes acne, or anaerobic bacteria.

15 9. The composition of claim 6, wherein the bacterium is a live bacterium, fragments of cell walls of the same, or culture filtrate of the same.

10. A cosmetic composition comprising at least one bacterium selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus fermentum*, and
20 a culture filtrate of bacteria thereof.

11. The cosmetic composition of claim 10, wherein the bacterium is at least one selected from the group consisting of *Lactobacillus*

coprophilus PL 9001 (KCCM-10245), *Enterococcus durans* PL 9002 (KCCM-10246), *Streptococcus faecalis* PL 9003 (KCCM-10247), *Lactobacillus coprophilus* PL 9004 (KCCM-10248), *Lactobacillus fermentum* PL 9005 (KCCM-10250), and *Lactobacillus fermentum* PL 9006 (KCCM-10251).

5 12. The cosmetic composition of claim 10, wherein the cosmetic composition is for external application against infectious disease.

 13. The cosmetic composition of claim 10, wherein the cosmetic composition is for treatment of acne.

 14. A food additive comprising at least one bacterium selected
10 from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

 15. The food additive of claim 14, wherein the food additive is applied to yogurt, weaning diet, dairy goods, cheese, Kimchi, drinks, or solid
15 food.

 16. A food prepared by fermenting at least one bacterium selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, and *Lactobacillus fermentum*.

 17. An immunological enhancement composition comprising at
20 least one bacterium selected from the group consisting of *Lactobacillus coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

18. The composition of claim 17, wherein the composition induces expression of tumor necrosis factor- α or interleukin-6.

19. A composition for curing intestinal disorders comprising at least one bacteria selected from the group consisting of *Lactobacillus* 5 *coprophilus*, *Enterococcus durans*, *Streptococcus faecalis*, *Lactobacillus fermentum*, and a culture filtrate of bacteria thereof.

20. The composition of claim 19, wherein the composition inhibits the growth of at least one bacterium selected from the group consisting of *Helicobacter pylori*, a bacillus that causes food poisoning, and 10 anaerobic bacteria.

1/32

Fig. 1

280 ug 140 ug 70 ug 14 ug 1.4 ug 0.14 ug

0ug

Fig. 2

Control	Competitive Lipid Binding Assay		
Lipid	PL9001	PL9002	PL9003
<i>Helicobacter pylori</i>	PL9004	PL9005	PL9006

2/32

Fig. 3a





Fig. 3b



3/32

Fig. 3c





Fig. 3d



4/32

Fig. 3e



Fig. 3f




5/32

Fig. 3g



Fig. 3h



6/32

Fig. 3i





Fig. 3j



7/32

Fig. 3k





Fig. 3l



8/32

Fig. 4a



(a)

(a')

Fig. 4b



(b)

(b')

9/32

Fig. 4c



(c)

(c')

Fig. 4d



(d)

(d')

2

10/32



Fig. 4e



(e)

(e')


Fig. 5



Diameter of Growth-Inhibited-zone


11/32

Fig. 6



12/32

Fig. 7



13/32

Fig. 8a





Fig. 8b



14/32

Fig. 8c





Fig. 8d



15/32

Fig. 8e





Fig. 9a



16/32

Fig. 9b





Fig. 9c



17/32

Fig. 9d





Fig. 9e



18/32

Fig. 10a





Fig. 10b



19/32

Fig. 10c





Fig. 10d



20/32

Fig. 10e





Fig. 11a



21/32

Fig. 11b





Fig. 11c



22/32

Fig. 11d





Fig. 11e



23/32

Fig. 12a





Fig. 12b



24/32

Fig. 12c





Fig. 12d



25/32

Fig. 12e





Fig. 13a



26/32

Fig. 13b





Fig. 13c



27/32

Fig. 13d





Fig. 13e



28/32

Fig. 14






Fig. 15a



29/32

Fig. 15b



30/32

Fig. 16a





Fig. 16b



31/32

Fig. 16c





Fig. 16d



32/32

Fig. 16e





Fig. 16f



Sequence Listing

<110> PL BIO Co., Ltd

<120> LACTIC ACID BACTERIA INHIBITING ADHESION OF HELICOBACTER PYLORI
TO GASTRIC MUCOSA

<130> opp20010966kr

<150> KR10-2000-74602

<151> 2000-12-08

<150> KR10-2001-2578

<151> 2001-01-17

<150> KR10-2001-2577

<151> 2001-01-17

<150> KR10-2001-2579

<151> 2001-01-17

<150> KR10-2001-8373

<151> 2001-02-20

<150> KR10-2001-40134

<151> 2001-07-05

<150> KR10-2001-40135

<151> 2001-07-05

<150> KR10-2001-40136

<151> 2001-07-05

<150> KR10-2001-40137

<151> 2000-07-05

<150> KR10-2001-40138

<151> 2001-07-05

<150> KR10-2001-40139

<151> 2001-07-05

<150> PCT/KR01/01286

<151> 2001-07-27

<150> KR10-2001-76748

<151> 2001-12-05

<150> KR10-2001-76749

<151> 2001-12-05

<150> KR10-2001-76750

<151> 2001-12-05

<150> KR10-2001-76751

<151> 2001-12-05

<150> KR10-2001-76752

<151> 2001-12-05

<160> 6

<170> KopatentIn 1.71

<210> 1

<211> 527

<212> DNA

<213> 16S rRNA sequence of Lactobacillus coprophilus PL 9001

<400> 1

cgctggcggc gtgcctaata catgcaagtc gaacgctttg tggttcaact gatttgaaga 60

gcttgctcag atatgacgat ggacattgca aagagtggcg aacgggtgag taacacgtgg 120

gaaacctacc tcttagcagg ggataacatt tggaacaga tgctaatacc gtataacaat 180

agcaaccgca tggttgctac ttaaaagatg gttctgctat cactaagaga tggccccg 240

gtgcattagt tagttgggta ggtaatggct caccaagacg atgatgcata gccgagttga 300

gagactgata ggccacaatg ggactgagac acggcccata ctctacggg aggcagcagt 360

agggaaatctt ccacaatggg cgaaagcctg atggagcaac gccgcgtgtg tgatgaaggg 420
 ttctggctcg taaaacactg ttgtaagaga agaatgacat tgagagtaac tgttcaatgt 480
 gtgacgggtat cttaccagaa aggaacggct aaatacgtgc cagcaag 527

<210> 2
 <211> 481
 <212> DNA
 <213> 16S rRNA sequence of Enterococcus durans PL 9002

<400> 2
 ggacaacctt gcggcgtgct aataatgcaa gtcgtacgct tctttttcca ccggagcttg 60
 ctccaccgga aaaagaagag tggcgaacgg gtgagtaaca cgtgggtaac ctgcccata 120
 gaaggggata acacttgga acaggtgcta ataccgtata acaatcgaaa ccgcatggtt 180
 ttgatttgaa aggcgctttc ggggtgcgct gatggatgga cccgcggtgc attagctagt 240
 tggtcaggta acggctcacc aaggccacta tgcatagccg acctgagagg gtgatcggcc 300
 acattgggac tgagacacgg cccaaactec tacgggaggg agcagtaggg aatcttcggc 360
 aatggacgaa agtctgaccg agcaacgccg cgtgagttaa gaaggttttc cgggaatcgt 420
 taaactctgt ttgttaagaa gaaagaacca aaggggattg aaggagttat cttgtttcat 480
 t 481

<210> 3
 <211> 825
 <212> DNA
 <213> 16S rRNA sequence of Enterococcus faecalis PL 9003

<400> 3
 ctggcggcgt gctaataatg caagcgaacg cttctttcct cccgagtgtg tgcaactcaat 60

tggaaagagg agtggcggac gggtagtaac cacgtgggta acctacccat cagaggggga 120
 taacacttgg aaacagggtgc taataccgca taacagtta tgccgcatgg cataagagtg 180
 aaaggcgctt tcgggtgtcg ctgatggatg gaccgcggt gcattagcta gttggtgagg 240
 taacggctca ccaaggccac gatgcatagc cgacctgaga gggtagtcgg ccacactggg 300
 actgagacac ggcccagact cctacgggag gcagcagtag ggaatcttcg gcaatggacg 360
 aaagtctgac cgagcaacgc cgcgttgagt gaaagaagggt ttttcgggat cgtaaaaaac 420
 tctgttgttt agaggaatct tcggatggac gaaagctgac cgagcaacgc cgcgtgagtg 480
 aagaaggttt tcggatcgta aaactctgtt gttagagaag aacaaggacg ttagtaactg 540
 aacgtccctt gacggtatct aaccagaaag ccacggctaa ctacgtgcca gcagccgagg 600
 taatacgtag gtggcaagcg ttgtccgat ttattgggag taaagcgagc gcaggcggtt 660
 tcttaagtct gatgtgaaag cccccggctc aaccggggag ggtcattgga aactgggaga 720
 cttgagtgca gaagaggaga gtggaattcc atgtgtagcg gtgaaatgag tagatatatg 780
 gaggaacacc agtggcgaag ggggtctctt ggtctgtaac tgacg 825

<210> 4

<211> 475

<212> DNA

<213> 16S rRNA sequence of *Lactobacillus coprophilus* PL 9004

<400> 4

acgtggcgg cgtgcetaat acatgcaagt cgaacgcttt gtgggtcaac tgatttgaag 60
 agcttgctca gatatgacga tggacattgc aaagagtggc gaacgggtga gtaacacgtg 120
 ggaaacctac ctcttagcag gggataacat ttggaaacag atgctaatac cgtataacaa 180
 tagcaaccgc atgggttgcta cttaaaagat ggttctgcta tcactaagag atgggtccgc 240

ggtgcattag ttagttggtg aggtaatggc tcaccaagac gatgatgcat agccgagttg 300
 agagactgat cggccacaat gggactgaga cacggcccat actcctacgg gaggcagcag 360
 tagggaatct tccacaatgg gcgaaagcct gatggagcaa cgccgcgtgt gtgatgaagg 420
 gtttcggctc gtaaaacact gttgtaagag aagaatgaca ttgagagtaa ctgtt 475

<210> 5

<211> 339

<212> DNA

<213> 16S rRNA sequence of Lactobacillus fermentum PL 9005

<400> 5

ggtgtgccta atacatgcaa gtcgaacgag ttggcccaat tgattgatgg tgcttgacc 60
 tgattgattt tggtcgcca cagatggcgg acgggtgagt aacacgtagg taacctgccc 120
 agaagcgggg gacaacattt ggaacagat gctaataccg cataacagcg ttgttcgcat 180
 gaacaacgct taaaagatgg cttctcgcta tcacttctgg atggacctgc ggtgcattag 240
 cttgttggtg gggtaacggc ctaccaaggc gatgatgcat agccgagttg agagactgat 300
 cggccacaat gggactgaga cacggcccat actcctacg 339

<210> 6

<211> 447

<212> DNA

<213> 16S rRNA sequence of Lactobacillus fermentum PL 9006

<400> 6

acgccggcgg tgtgccta atcatgcaagt cgaacgcgtt ggcccaattg attgatggtg 60
 cttgcacctg attgattttg gtcgccaacg agtggcggac gggtagta cacgtaggta 120
 acctgcccag aagcggggga caacatttgg aaacagatgc taataccgca taacagcgtt 180

gttcgcatga acaacgctta aaagatggct tctcgctatc acttctggat ggacctgcgg	240
tgcattagct tgttggtggg gtaanggcct accaaggcga tgatgcatag ccgagttgag	300
agactgatcg gccacantgg gactgagaca cggcccatac tcctacggga ggcagcagta	360
gggaatcttc cacaatgggc gcaagcctga tggagcaaca ccgcgtgagt gaagaagggt	420
ttcggtcgt aaagctctgt tgttaaa	447

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR 01/02126

CLASSIFICATION OF SUBJECT MATTER

IPC⁷: A61K 35/74, C12N 1/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁷: A61K 35/74

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CAS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/55131 A1 (GRAHN HAKANSSON, E. et al.) 10 December 1998 (10.12.98) <i>page 3, lines 6-14; page 4, lines 19-36; page 6, lines 17-30; page 7, lines 3-25; page 9, lines 8-12.</i>	1,3,4,6,8,9, 14-17,19,20
X	EP 1000625 A1 (LABORATOIRE DU LACTEOL DU DOCTEUR BOUCARD, INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE) 17 May 2000 (17.05.00) <i>abstract; page 1, lines 25-27; claims 1,3,4,6.</i>	1,3,4,6,8,9
X	EP 1038951 A1 (SOCIETE DES PRODUITS NESTLE S.A.) 27 September 2000 (27.09.00) <i>page 1, lines 6-10, 51-54.</i>	1,3,4,6,8,9, 14-16,19,20
X	JP 09 241173 A (WAKAMOTO PHARM CO LTD) 16 September 1997 (16.09.97) (abstract) WPI [online]. London, U.K.: Derwent Publications, Ltd. [retrieved on 13-02-2002]. DW 199750, AN: 1997-539731 [50] <i>abstract.</i>	1,3,4,6,8,9, 14-16,19,20

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

„A“ document defining the general state of the art which is not considered to be of particular relevance.

„E“ earlier application or patent but published on or after the international filing date

„L“ document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

„O“ document referring to an oral disclosure, use, exhibition or other means

„P“ document published prior to the international filing date but later than the priority date claimed

„T“ later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

„X“ document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

„Y“ document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

„&“ document member of the same patent family

Date of the actual completion of the international search

14 February 2002 (14.02.2002)

Date of mailing of the international search report

14 March 2002 (14.03.2002)

Name and mailing address of the ISA/AT

Austrian Patent Office
Kohlmarkt 8-10; A-1014 Vienna
Facsimile No. 1/53424/535

Authorized officer

MOSSER

Telephone No. 1/53424/437

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 01/02126

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 63 179829 A (ADVANCE KK) 23 July 1988 (23.07.88) (abstract) WPI [online]. London, U.K.: Derwent Publications, Ltd. [retrieved on 13-02-2002]. DW198835, AN: 1988-246702 [25] <i>abstract.</i>	10,12,13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR 01/02126

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
EP	A1	1000625	17-05-2000	FR	A1 2785190	05-05-2000
				FR	B1 2785190	26-01-2001
EP	A1	1038951	27-09-2000	BR	A 00001392	02-05-2001
				JP	A2 00279166	10-10-2000
				US	BA 6340585	22-01-2002
JP	A2	9241173	16-09-1997		none	
JP	A2	63179829	23-07-1988	JP	B4 8018995	28-02-1996
WO	A1	9855131	10-12-1998	AU	A1 80460/98	21-12-1998
				AU	B2 726217	02-11-2000
				CN	T 1272791	08-11-2000
				EP	A1 1005353	07-06-2000
				NO	A0 995794	26-11-1999
				NO	A 995794	31-01-2000
				SE	A0 9702083	02-06-1997
				SE	A 9702083	03-12-1998
				SE	C2 511524	11-10-1999

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.